arXiv:1201.1727v1 [nlin.CD] 9 Jan 2012

Relaxation dynamics of the Kuramoto model with uniformly distributed natural frequencies

Anandamohan Ghosh[1] and Shamik Gupta[2]
[1]*Indian Institute of Science Education and Research-Kolkata, Mohanpur 741252, India*
[2]*Laboratoire de Physique de l'École Normale Supérieure de Lyon,*
Université de Lyon, CNRS, 46 Allée d'Italie, 69364 Lyon cédex 07, France

(Dated: January 10, 2012)

The Kuramoto model describes a system of globally coupled phase-only oscillators with different natural frequencies. The model in the steady state exhibits a phase transition as a function of the coupling parameter between a low-coupling incoherent phase in which the oscillators oscillate independently and a high-coupling synchronized phase in which they oscillate with a common frequency. Here, we consider a uniform distribution for the natural frequencies, for which the phase transition is of first oder. We study how the system close to the phase transition relaxes in time to the steady state while starting from an initial homogeneous state. In this case, the relaxation process occurs as a step-like jump in the order parameter, and has been previously interpreted as suggesting the existence of metastable states. By analyzing (i) the distribution of the jump time, and (ii) the distribution of the order parameter in the steady state, we provide evidence to rule out the existence of any metastable states.

PACS numbers: 05.45.Xt

Systems of coupled oscillators of different natural frequencies often exhibit collective synchronization in which a finite fraction of the oscillators oscillates with a common frequency. Examples include groups of fireflies flashing in unison [1, 2], networks of pacemaker cells in the heart [3, 4], superconducting Josephson junctions [5, 6], and many others. Understanding the nature and emergence of synchronization from the underlying dynamics of such systems is an issue of great interest. A paradigmatic model in this area is the so-called Kuramoto model involving globally-coupled oscillators [7]. Although studied extensively in the past, the model continues to raise new questions, and has been a subject of active research; for reviews, see [8, 9].

The model consists of N phase-only oscillators labeled by the index $i = 1, 2, \ldots, N$. Each oscillator has its own natural frequency ω_i distributed according to a given probability density $g(\omega)$, and is coupled to all the other oscillators. The phase of the oscillators evolves in time according to [7]

$$\frac{d\theta_i}{dt} = \omega_i + \frac{K}{N}\sum_{j=1}^{N}\sin(\theta_j - \theta_i), \qquad (1)$$

where θ_i, the phase of the ith oscillator, is a periodic variable of period 2π, and $K \geq 0$ is the coupling constant.

The Kuramoto model has been mostly studied for a unimodal $g(\omega)$, i.e., one which is symmetric about the mean frequency $\omega = \Omega$, and which decreases monotonically and continuously to zero with increasing $|\omega - \Omega|$ [8, 9]. Then, it is known that in the limit $N \to \infty$, the system of oscillators in the steady state undergoes a continuous transition at the critical threshold $K_c = 2/\pi g(0)$. For $K < K_c$, each oscillator tends to oscillate independently with its own natural frequency. On the other hand, when $K > K_c$, the coupling synchronizes the phases of the oscillators, and in the limit $K \to \infty$, they all oscillate with the mean frequency Ω.

The degree of synchronization in the system at time t is measured by the complex order parameter

$$\mathbf{r}(t) = r(t)e^{i\psi(t)} = \frac{1}{N}\sum_{j=1}^{N} e^{i\theta_j(t)}, \qquad (2)$$

with magnitude $r(t)$ and phase $\psi(t)$, in terms of which the time evolution (1) may be written as

$$\frac{d\theta_i}{dt} = \omega_i + Kr(t)\sin(\psi(t) - \theta_i). \qquad (3)$$

Here $r(t)$ with $0 \leq r(t) \leq 1$ measures the phase coherence of the oscillators, while $\psi(t)$ gives the average phase. When K is smaller than K_c, it is known that $r(t)$ while starting from any initial value relaxes in the long-time limit to zero, corresponding to an incoherent phase in the steady state. For $K > K_c$, on the other hand, $r(t)$ grows in time to asymptotically saturate to a non-zero value $r(K) \leq 1$ that increases continuously with K.

For a unimodal $g(\omega)$, the relaxation of $r(t)$ to steady-state is exponentially fast for $K > K_c$. For $K < K_c$, however, the nature of relaxation depends on $g(\omega)$. When $g(\omega)$ has a compact support, $r(t)$ while starting from any initial value decays to zero more slowly than any exponential as $t \to \infty$ [10]. When $g(\omega)$ is supported on the whole real line, $r(t)$ as a function of time is known only in particular cases. For example, for a Lorentzian $g(\omega)$, and a specific initial condition, $r(t)$ decays exponentially to zero [10]. For other choices of $g(\omega)$ in this class and for other initial conditions, the dependence of $r(t)$ on time is not known analytically, and it has been speculated that $r(t)$ is a sum of decaying exponentials [10].

In the limit $N \to \infty$, the state of the oscillator system at time t is described by the probability distribution $f(\theta, t, \omega)$ that gives for each natural frequency ω

the fraction of oscillators with phases between θ and $\theta + d\theta$ at time t. The time evolution of $f(\theta, t, \omega)$ satisfies the continuity equation for the conservation of the number of oscillators with natural frequency ω, and is given by a non-linear partial integro-differential equation [8]. Recent analytical studies for a unimodal $g(\omega)$ (specifically, a Lorentzian) and for a bimodal $g(\omega)$ (sum of two Lorentzians) demonstrated by considering a restricted class of $f(\theta, t, \omega)$ that the time evolution in terms of the integro-differential equation may be exactly reduced to that of a small number of ordinary differential equations (ODEs) [11, 12]. Interestingly, this system of ODEs for the reduced system contains the whole spectrum of dynamical behavior of the full system. However, the validity of such a reduction for a general $g(\omega)$ is not known.

A uniform $g(\omega)$ with a compact support does not qualify as a unimodal distribution. In this case, it is known that in the limit $N \to \infty$, the Kuramoto model in the steady state exhibits a first-order phase transition between an incoherent and a synchronized phase at the critical coupling $K_c = 2/\pi g(0)$ [13]. For finite N, study of the relaxation of an initial state with uniformly distributed phases has demonstrated that for values of K around K_c, the process occurs as a step-like jump in $r(t)$ from its initial to the steady state value. This behavior has been interpreted as suggesting the existence of metastable states in the system [14, 15].

In this work, we investigate in detail the implication of the existence of the step-like relaxation. By performing extensive numerical simulations, we analyze in detail the distribution $P(\tau)$ of the jump time τ, and also the distribution $P(r)$ of the steady state value r of the order parameter. Our analysis suggests the absence of any metastable states in the system.

The numerical simulations involved integrating Eq. (3) by a 4th-order Runge-Kutta algorithm. In simulations, we monitored the time evolution of $r(t)$ while starting from an initial state with θ_i's independently and uniformly distributed in $[-\pi, \pi]$ and ω_i's independently and uniformly distributed in $[-2, 2]$ (so that $g(\omega) = 1/4$). Thus, initially, $r(t) \approx 0$. We call such an initial state a homogeneous state.

For $K = 2.62$, close to the critical value $K_c = 2/\pi g(0) \approx 2.55$, Fig. 1(a) shows the temporal evolution of $r(t)$ for 8 different realizations of the initial homogeneous state for $N = 1000$, showing that $r(t)$ jumps in a step at times ~ 100 from its initial to its steady state value ≈ 0.82 (with fluctuations). For the realization shown in Fig. 1(b), however, the jump to the steady state value takes place at time ≈ 2000, while for the realization in Fig. 1(c), $r(t)$ stays close to the initial value, and the jump does not take place in the time duration shown. The average of $r(t)$ over the realizations in (a)-(c) is shown in Fig. 1(d), which naturally exhibits a step-like relaxation. Similar behavior of the average $r(t)$, where the averaging is over a *few* realizations, was reported in [14]. Figure 1(e) shows the time evolution of $r(t)$ when averaged over a *large* (~ 1000) number of realizations, which however does not show any step-like relaxation.



FIG. 1: (a) Evolution of $r(t)$ in time for 8 different realizations of the initial homogeneous state. Here, $K = 2.62$, while the critical coupling is $K_c \approx 2.55$. The plot shows that for these realizations, $r(t)$ jumps in a step at times ~ 100 from its initial to its steady state value. (b) and (c) also show the time evolution of $r(t)$ for two other realizations of the initial state: for the realization in (b), $r(t)$ shows a jump to the steady state value at time ≈ 2000, while for the one in (c), $r(t)$ remains close to its initial value and does not relax to the steady state value in the time shown. (d) and (e) show respectively the time evolution of $r(t)$ when averaged over the realizations in (a)-(c), and over typical 1000 realizations. The data are obtained from numerical integration of the N-body dynamics (3) with $N = 1000$.

Figure 1 illustrates that the relaxation of $r(t)$ occurs as a step-like jump, and that there is a wide distribution of the jump time τ. Thus, it is worthwhile to study the distribution $P(\tau)$. Our numerical data, displayed in Fig. 2, show that $P(\tau)$ has a power-law tail; the scaling collapse of the data, shown in Fig. 3, suggests the following scaling form:

$$P(\tau) \sim \sqrt{N} f\left(\frac{\tau}{\sqrt{N}}\right), \tag{4}$$

where $f(x)$ is the scaling function.

In the power-law regime of $P(\tau)$, let us write

$$P(\tau) \propto \tau^{-\alpha}, \tag{5}$$

where α is the decay exponent. Then, in this long-time regime, $r(t)$ at time t will have a value close to one provided that the jump from the initial close-to-zero value has already taken place by this time; otherwise, $r(t) \approx 0$. It then follows that the average order parameter, $\langle r \rangle(t)$,



FIG. 2: Distribution $P(\tau)$ of the time τ at which $r(t)$ jumps from a close-to-zero value corresponding to an initial homogeneous state to the final steady state value (see Fig. 1(b) for an example of such a jump). The data are obtained from numerical integration of the N-body dynamics (3). Here, $K = 2.62$. The inset shows power-law fits for the tail of $P(\tau)$ according to Eq. (5), giving values 2.2, 2.7, and 2.8 for system sizes $N = 1000$, 2000 and 4000, respectively.



FIG. 3: Scaling collapse of the data for the distribution $P(\tau)$ shown in Fig. 2.

in this regime will be

$$\langle r\rangle(t) \sim \int_0^t d\tau\ P(\tau) \sim 1 - t^{1-\alpha}. \tag{6}$$

Equivalently, one has

$$1 - \langle r\rangle(t) \sim t^{-\beta}, \tag{7}$$

with

$$\beta = \alpha - 1. \tag{8}$$



FIG. 4: (a) Power-law fit to the tail of $P(\tau)$, according to Eq. (5). (b) Power-law fit to the quantity $1 - \langle r\rangle(t)$, according to Eq. (7). The data are obtained from numerical integration of the N-body dynamics (3) with $N = 1000$, and for $K = 2.62$. The average is computed over ~ 1000 realizations.

This prediction is easily checked from our simulation results. Figure 4(b) shows the power-law fit to the tail of $P(\tau)$ for $N = 1000$, giving $\alpha \approx 2.2$. On the other hand, Fig. 4(b) shows the power-law fit to the quantity $1 - \langle r\rangle(t)$ at long times, giving $\beta \approx 0.9$ in Eq. (7), while the value predicted by Eq. (8) is 1.2. One may hope to obtain a closer agreement with better statistics.

To probe the existence of metastable states in the system, we obtained the probability distribution of the steady state value r of the order parameter. Figure 5 shows that with increasing system size, the distribution is peaked at a value close to one, while the probability to obtain any other value is negligibly small. We have checked that the distribution $P(r)$ shown in the figure does not evolve in time, and thus, corresponds to the steady state. The form of $P(r)$ confirms the absence of metastable states in the thermodynamic limit, whose presence would have otherwise led to peaks in $P(r)$ at values close to zero.

Based on the above analysis, we thus conclude that

1. For uniformly distributed frequencies and a homogeneous initial state, the order parameter in the Kuramoto model shows relaxation as a step-like jump from its initial value (≈ 0) to its steady state value (≈ 1) for values of K close to the phase transition ($K \gtrsim K_c$).

2. There is a wide distribution of the time τ of occurrence of the jump, and its distribution $P(\tau)$ has a power-law tail.

3. Hence, averaging $r(t)$ over a few realizations naturally shows a step-like relaxation, and may erroneously lead to the conclusion that the system during the relaxation gets stuck in metastable states.

4. In order to capture the whole range of values of τ as suggested by its fat-tailed distribution, it is thus important that $r(t)$ is averaged over a very large number of realizations. We demonstrated that in



FIG. 5: Starting from an initial homogeneous state, the figure shows the distribution $P(r)$ of the steady state value r of the order parameter. The data are obtained from numerical integration of the N-body dynamics (3) for the values of N shown in the figure, and for $K = 2.62$.

doing so, $r(t)$ does not show any step-like relaxation, but instead a smooth evolution towards the steady state. This is consistent with the absence of any metastable states in the system, and is further corroborated by our results on the distribution $P(r)$ of the steady state order parameter, which is sharply peaked around a value close to 1 in the thermodynamic limit.

We obtained a scaling form for the distribution $P(\tau)$. We also predicted and verified numerically a power-law growth of the averaged order parameter at late times. However, we do not have an understanding of the relaxation mechanism, and an investigation in that direction is in progress.

Acknowledgments

S. G. acknowledges support of the French contract LORIS (ANR-10-CEXC-010-01). We thank Deepak Dhar, David Mukamel, Antonio Politi, and Stefano Ruffo for useful discussions and comments on the manuscript.

[1] J. Buck and E. Buck, Sci. Am. **234**, 74 (1976).
[2] J. Buck, Quart. Rev. Biol. **63**, 265 (1988).
[3] C. S. Peskin, Math. Aspects of Heart Physiology, Courant Institute of Mathematical Science Publications, New York, 1975, pp. 268-278.
[4] D. C. Michaels, E. P. Matyas, and J. Jalife, Circulations Res. **61**, 704 (1987).
[5] K. Wiesenfeld, P. Colet, and S. H. Strogatz, Phys. Rev. Lett. **76**, 404 (1996).
[6] K. Wiesenfeld, P. Colet, and S. H. Strogatz, Phys. Rev. E **57**, 1563 (1998).
[7] Y. Kuramoto, in *International Symposium of Mathematical Problems in Theoretical Physics*, edited by H. Araki, Lecture Notes in Physics Vol. 39 (Springer, Berlin, 1975).
[8] S. H. Strogatz, Physica D **143**, 1 (2000).
[9] J. A. Acebrón, L. L. Bonilla, C. J. Pérez Vicente, F. Ritort, and R. Spigler, Rev. Mod. Phys. **77**, 137 (2005).
[10] S. H. Strogatz, R. E. Mirollo, and P. C. Matthews, Phys. Rev. Lett. **68**, 2730 (1992).
[11] E. Ott and T. A. Antonsen, Chaos **18**, 037113 (2008).
[12] E. A. Martens, E. Barreto, S. H. Strogatz, E. Ott, P. So, and T. M. Antonsen, Phys. Rev. E **79**, 026204 (2009).
[13] D. Pazó, Phys. Rev. E **72**, 046211 (2005); L. Basnarkov and V. Urumov, Phys. Rev. E **76**, 057201 (2007); Phys. Rev. E **78**, 011113 (2008).
[14] A. Pluchino and A. Rapisarda, Physica A **365** 184 (2006).
[15] G. Miritello, A. Pluchino, and A. Rapisarda, Europhys. Lett. **85**, 10007 (2009).